

ABGENEX Pvt. Ltd., E-5, Infocity, KIIT Post Office, Tel: +91-674-2720712, +91-9437550560 Email: info@abgenex.com Bhubaneswar, Odisha - 751024, INDIA

32-3149: A2M Recombinant Protein

Alternative Name: Alpha-2-macroglobulin, Alpha-2-M, A2M, CPAMD5, FWP007, S863-7, alpha 2M, DKFZp779B086.

Description

Source: Human Plasma. Human Alpha-2 Macroglobulin is a tetrameric glycoprotein, produced in Human plasma and having a molecular mass of 725 kDa. Alpha-2 Macroglobulin is a serine proteases inhibitor, which inhibits coagulation by inactivating thrombin and Kallikrein, it inhibits fibrinolysis by inactivating plasmin and involved in insulin transport. Alpha-2-Macroglobulin is a large plasma protein, which is produced by the liver, it's composed of 4 identical subunits bound together by -S-S- bonds. A2M is able to inactivate many kinds of proteinases (including serine-, cysteine-, aspartic- and metalloproteinases). A2M has a 35 amino acid 'bait' region in its structure. Proteinases bind and cleave the 'bait' region become bound to A2M. Macrophage receptors recognize the proteinase-A2M complex and clear it from the system. A2M binds to and removes MMP-2 and MMP-9 (active forms of the gelatinase) from the circulation using scavenger receptors on the phagocytes. The levels of Alpha-2-macroglobulin are increased in nephrotic syndrome which is a condition where the kidneys start to leak out some of the smaller blood proteins. Due to its large size, A2-macroglobulin is retained in the bloodstream. Increased production of all proteins causes A2-macroglobulin concentration to increase. Chronic renal failure might lead to amyloid by alpha-2-macroglobulin. A2M is raised in cirrhosis, pregnancy and diabetes.

Product Info

Amount: 1 ma

Purification: Greater than 96.0%.

Content: Lyophilized from 0.02M TRIS, 0.15M NaCl and 0.1M Sucrose, pH 8.0.

Storage condition:

Human Alpha-2 Macroglobulin although stable at room temperature for 3 weeks, should be stored

between 2-8°C. Do not freeze!

Application Note

It is recommended to reconstitute the lyophilized A2M in sterile 18M Ω -cm H2O.

