

ABGENEX Pvt. Ltd., E-5, Infocity, KIIT Post Office, Tel: +91-674-2720712, +91-9437550560 Email: info@abgenex.com

Bhubaneswar, Odisha - 751024, INDIA

## 32-6120: Mouse Anti Human Immature Colon Carcinoma Transcript 1(Clone:PAT1E9A)

Clone Name: Monoclonal
Clone Name: PAT1E9A
Application: ELISA,WB
Gene: ICT1

 Gene :
 ICT1

 Gene ID :
 3396

 Uniprot ID :
 Q14197

 Format :
 Purified

Alternative Name:

DS-1,DS1,Peptidyl-tRNA hydrolase ICT1 mitochondrial,Digestion substraction 1,Immature colon

carcinoma transcript 1 protein,ICT1.

**Isotype:** Mouse IgG1 heavy chain and Kappa light chain.

Anti-human ICT1 mAb, clone PAT1E9A, is derived from hybridization of mouse F0 myeloma cells

Immunogen Information: with spleen cells from BALB/c mice immunized with a recombinant human ICT1 protein 30-206

amino acids purified from E. coli.

## **Description**

Immature Colon Carcinoma Transcript 1 (ICT1) functions as a codon-independent translation release factor that has lost all stop codon specificity and leads the termination of translation in mitochondrion in case of abortive elongation. The mature colon epithelium has 3 differentiated cell types which arise from a multipotent stem cell. ICT1 is involved in the hydrolysis of peptidyltRNAs which have been prematurely terminated and consequently in the recycling of stalled mitochondrial ribosomes. Digression from the normal maturation pathway by neoplastic transformation is assumed to originate in stem cells or their early descendants.

## **Product Info**

Amount: 20 µg

**Purification:** ICT1 antibody was purified from mouse ascitic fluids by protein-A affinity chromatography.

Content: 1mg/ml containing PBS, pH-7.4, 10% Glycerol and 0.02% Sodium Azide.

Storage condition:

For periods up to 1 month store at 4°C, for longer periods of time, store at -20°C. Prevent freeze

thaw cycles.

## **Application Note**

The antibody has been tested by ELISA, Western blot analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results. Recommended starting dilution is 1:1000.