

## 10-6006: Monoclonal Antibody to RANKL (Clone: ABM10A7)

| Clonality :             | Monoclonal   |
|-------------------------|--|
| Clone Name :            | ABM10A7  |
| Application :           | FACS,WB  |
| Reactivity :            | Mouse,Human  |
| Gene :                  | TNFSF11  |
| Gene ID :               | 8600   |
| Uniprot ID :            | O14788   |
| Format :                | Purified   |
| Alternative Name :      | Tumor necrosis factor ligand superfamily member 11,Osteoclast differentiation factor,Osteoprotegerin ligand,Receptor activator of nuclear factor kappa-B ligand,TNF-related activation-induced cytokine,TRANCE,CD254 |
| Isotype :               | Mouse IgG1, Kappa  |
| Immunogen Information : | Full length recombinant protein of RANKL was used as the immunogen for this antibody.  |

## **Product Info**

| Amount :            | 25 μg / 100 μg   |
|---------------------|--|
| Purification :      | Protein G Chromatography   |
| Content :           | 25 $\mu g$ in 50 $\mu l/100~\mu g$ in 200 $\mu l$ PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic. |
| Storage condition : | Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.    |

## **Application Note**

WB: 2-4 µg/ml, FACS: 0.5-1 µg/10^6



Fig-1: Western blot analysis of RANKL. Anti-RANKL antibody (Clone: ABM10A7) was used at 2  $\mu$ g/ml on mPlacenta tissue lysate.





Fig-2: Intra cellular flow analysis of RANKL in Jurkat using 0.5 µg/10^6 cells of antibody (Clone: ABM10A7). Green represents isotype control; red represents anti-RANKL antibody. Goat anti-mouse PE conjugate was used as secondary antibody.

Fig-3: Intra cellular flow analysis of RANKL in HepG2 using  $0.5 \mu g/10^{6}$  cells of antibody (Clone: ABM10A7). Green represents isotype control; red represents anti-RANKL antibody. Goat anti-mouse PE conjugate was used as secondary antibody.