

10-3012-F: Monoclonal antibody to MyD88 (Clone: ABM2H20)

Clonality :	Monoclonal
Clone Name :	ABM2H20
Application :	FACS
Reactivity :	Human
Conjugate :	FITC
Gene :	MYD88
Gene ID :	4615
Uniprot ID :	Q99836
Format :	Purified
Isotype :	Mouse IgG1 Kappa
Immunogen Information :	A partial length recombinant protein of human MyD88 (amino acids 13-221) was used as an immunogen for this antibody.

Description

MyD88 (Myeloid differentiation factor) is an essential adaptor molecule in all TLR (Toll-like receptor) signaling pathways except TLR3. MyD88 is composed of an N-terminal (Death Domain) and a highly conserved C-terminal TIR (Toll/interleukin-1 Receptor) domain. It is found to stimulate IL-1R/IL18R-mediated signaling. MyD88-dependent signaling is also important in the regulation of innate as well as acquired immunity, in particular, T-cell responses, to various microbial pathogens. After activation of TLRs, MyD88 is phosphorylated and subsequently recruits IRAKs (IL-1R Associated Kinases) and other downstream proteins such as TRAF6, finally resulting in activation of the NF-kappaB (nuclear factor kappa B) pathway.

Product Info

Amount :	25 µg / 100 µg
Purification :	Protein G Chromatography
Content :	0.2 mg/ ml of Tris buffer containing 0.05% sodium azide.
Storage condition :	Store the antibody at 4°C; stable for 6 months.

Application Note

FACS Analysis: 0.5-1 µg/ml




Fig:1-Intracellular flow analysis of MyD88 in PBMC using 0.5 µg/10⁶ cells. Green represents isotype control (ABEOMICS); red represents FITC conjugated anti-MyD88 antibody (10-3012F).