

10-7618: Monoclonal Antibody to Galectin-1 (Clone: ABM5B54)

Clonality :	Monoclonal
Clone Name :	ABM5B54
Application :	IHC,FACS,WB
Reactivity :	Mouse,Human
Gene :	LGALS1
Gene ID :	3956
Uniprot ID :	P09382
Format :	Purified
Alternative Name :	14 kDa laminin-binding protein, 14 kDa lectin, Beta-galactoside-binding lectin L-14-1, Galaptin, HBL, HPL, Lactose-binding lectin 1, Lectin galactoside-binding soluble 1, Putative MAPK-activating protein PM12, S-Lac lectin 1
Isotype :	Mouse IgG2b, Kappa
Immunogen Information	: Full length recombinant Galectin-1 protein was used as the immunogen for this antibody.

Product Info

Amount :	25 µg / 100 µg
Purification :	Protein G Chromatography
Content :	25 μg in 50 $\mu l/100~\mu g$ in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles

Application Note

Western blot analysis: 2-4 µg/ml, FACS analysis-0.5-1 µg/10^6 Cells, Immunohistochemical analysis-2-4 µg/ml




Figure:1- Western blot analysis of Galectin-1. Anti-Galectin-1 antibody (Clone: ABM5B54) was tested at 0.01 μ g/ml on (1) Recombinant protein and 2 μ g/ml on (2) Hela, (3) 3T3, (4) PC3, (5) 293, (6) THP1, (7) K562 and (8) HCT-116 lysates.






Figure:2- Intracellular flow analysis of Galectin-1 in U87 cell line using 0.5 µg/10^6 cells of Galectin-1 antibody (Clone: ABM5B54). Green represents isotype control; red represents anti-Galectin-1 antibody (10-7618). Goat anti-mouse PE conjugate was used as secondary antibody.

Figure:3- Intracellular flow analysis of Galectin-1 in A431 cell line using 0.5 μ g/10^6 cells of Galectin-1 antibody (Clone: ABM5B54). Green represents isotype control; red represents anti-Galectin-1 antibody (10-7618). Goat anti-mouse PE conjugate was used as secondary antibody.

Figure:4-Immunohistochemical analysis of Galectin-1. Anti-Galectin-1 (Clone: ABM5B54) was used in Colon Carcinoma tissue at 2 μ g/ml.