

### 36-1661: Monoclonal Antibody to S100A1 (Astrocyte Marker)(Clone : 4C4.9)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	4C4.9
<b>Application :</b>	IF,IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	S100A1
<b>Gene ID :</b>	6271
<b>Uniprot ID :</b>	P23297
<b>Format :</b>	Purified
<b>Alternative Name :</b>	S100A1,S100A
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Purified bovine brain S100 protein

#### Description


S100 belongs to the family of calcium binding proteins. S100A and S100B proteins are two members of the S100 family. S100A is composed of an alpha and beta chain whereas S100B is composed of two beta chains. S-100 protein has been found in normal melanocytes, Langerhans cells, histiocytes, chondrocytes, lipocytes, skeletal and cardiac muscle, Schwann cells, epithelial and myoepithelial cells of the breast, salivary and sweat glands, as well as in glial cells. Neoplasms derived from these cells also express S-100 protein, albeit non-uniformly. A large number of well-differentiated tumors of the salivary gland, adipose and cartilaginous tissue, and Schwann cell-derived tumors express S-100 protein. Almost all malignant melanomas and cases of histiocytosis X are positive for S-100 protein. Despite the fact that S-100 protein is an ubiquitous substance, its demonstration is of great value in the identification of several neoplasms, particularly melanomas and their metastases.

#### Product Info


<b>Amount :</b>	100 µg
<b>Purification :</b>	Affinity Chromatography
<b>Content :</b>	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

#### Application Note


Immunofluorescence (1-2ug/ml); ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),




Formalin--paraffin human Melanoma stained with S100A1 Monoclonal Antibody (4C4.9) (AEC Chromogen)



Western Blot of human Cerebellum Lysate using S100A1 Monoclonal Antibody (4C4.9).



Confocal Immunofluorescent analysis of A2058 cells using AF488-labeled S100A1 Monoclonal Antibody (4C4.9) (Green).



Confocal Immunofluorescent analysis of A2058 cells using AF488-labeled Isotype Control Monoclonal Antibody (IgG2a) (Green). F-actin filaments were labeled with DyLight 554 Phalloidin (red). DAPI was used to stain the cell nuclei (blue). (Negative Control)