

ABGENEX Pvt. Ltd., E-5, Infocity, KIIT Post Office, Tel: +91-674-2720712, +91-9437550560 Email: info@abgenex.com Bhubaneswar, Odisha - 751024, INDIA

32-1063: rBD 1 Recombinant Protein

Alternative Name: Beta-defensin 1,BD-1,rBD-1,Defensin beta 1,Defb1.

Description

Source: Escherichia Coli. BD-1 Rat Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing 37 amino acids and having a molecular mass of 4.1kDa. The BD-1 is purified by proprietary chromatographic techniques. The Defensin family are highly similar in their protein sequence and are microbicidal & cytotoxic peptides made by neutrophils. Beta Defensin-1 is an antimicrobial peptide having the resistance of epithelial surfaces to microbial colonization. Beta Defensin-1 has close proximity to Defensin Alpha-1 and has been implicated in the pathogenesis of cystic fibrosis. Skin of patients having atopic dermatitis patients and mycosis fungoides (non-lesional and lesional) show lower human Beta Defensin-1 mRNA expression and higher human Beta Defensin-2 and human Beta Defensin-3 mRNA expression. Beta Defensin is highly expressed by epithelial cells. Beta-defensin 1 may play a role in the pathogenesis of severe sepsis.

Product Info

Amount: 20 µg

Purification: Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: BD-1 protein was lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.

Lyophilized BD-1 although stable at room temperature for 3 weeks, should be stored desiccated

Storage condition: below -18°C. Upon reconstitution BD-1 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or

BSA). Please prevent freeze-thaw cycles.

Amino Acid: DQYRCLQNGG FCLRSSCPSH TKLQGTCKPD KPNCCRS.

Application Note

It is recommended to reconstitute the lyophilized BD-1 in sterile 18M-cm H2O not less than $100\text{\AA}\mu\text{g/ml}$, which can then be further diluted to other aqueous solutions. Measured by its ability to chemoattract CD34+ dendritic cells using a concentration range of 0.1-1.0 ug/ml.

