

## 32-1425: mL 7 Recombinant Protein

**Alternative Name :** Lymphopoietin 1 (LP-1),pre-B cell factor,IL-7.

### Description

Source : Escherichia Coli. Interleukin-7 Mouse Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 130 amino acids and having a molecular mass of 15.0 kDa. The IL-7 is purified by proprietary chromatographic techniques. Interleukin-7 is a potent lymphoid cell growth factor produced primarily by stromal cells. IL-7 has been shown to support the proliferation and differentiation of pre B- and early T cells as well as displaying a biological effect on cells of NK and myeloid lineages.

### Product Info

<b>Amount :</b>	10 µg
<b>Purification :</b>	Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
<b>Content :</b>	Filtered (0.2µm) and lyophilized from a concentrated (1mg/ml) solution in 1xPBS, pH7.4.
<b>Storage condition :</b>	Lyophilized Interleukin-7 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL7 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
<b>Amino Acid :</b>	ECHIKDKEGK AYESVLMISI DELDKMTGTD SNCPNNEPNF FRKHVCDDTK EAAFLNRAAR KLLKQFLKMNI SEEFNVHLLT VSQGTQTLVN CTSKEEKNVK EQKKNACFL KRLLEIKTC WNLKILKGS.

### Application Note

It is recommended to reconstitute the lyophilized Interleukin 7 in sterile 18M<sup>Ω</sup>-cm H<sub>2</sub>O not less than 100<sup>Å</sup>µg/ml, which can then be further diluted to other aqueous solutions. The ED<sub>50</sub> was determined by the dose-dependent stimulation of the proliferation of murine 2E8 cells is less than 0.2 ng/ml, corresponding to a specific activity of >5.0<sup>Å</sup>—106 IU/mg.

