

ABGENEX Pvt. Ltd., E-5, Infocity, KIIT Post Office, Tel: +91-674-2720712, +91-9437550560 Email: info@abgenex.com Bhubaneswar, Odisha - 751024, INDIA

16-1001: Quick-blots (Mouse Tissues)

Application : WB Reactivity : Mouse

# **Description**

**Description:** Quick-blots<sup>™</sup>

is Ready-to-use PVDF membrane with 11 mouse tissue lysates (20 µg total protein per lane). Quick-blots<sup>™</sup> is a very simple method to identify different mouse proteins.

## **Product Info**

Amount: 1 Membrane

**Content:** 1 pre-transferred membrane

**Storage condition :** Store unopened at RT. For long-term, store at 4?C.

## **Application Note**

#### Quick-blots<sup>™</sup>

# Protocol:

- 1. Activate the membrane with Methanol for 1-2 min and then wash the blots with TBST (25 mM Tris-Cl, pH 8.0; 125 mM NaCl; 0.1% Tween 20) buffer thrice to get rid of extra methanol.
- 2. Block the membrane for 1 hour in 5% skimmed milk (in TBST) to avoid non specific binding of antibodies.
- 3. Wash the membranes 2 times in TBST and incubate with primary antibodies prepared in 1% skimmed milk for 2 hr in RT/ over night at 4â•°C.
- 4. After primary incubation, wash the membranes for 1 hour in TBST at RT. (Buffer should be changed in 30 min interval)
- 5. Incubate the membranes in appopriate Secondry antibody for 1 hour.
- 6. After secondary incubation, wash the membranes for 2 hours (Buffer should be changed in 30 min interval). Then develop the blots with Chemiluminescent Substrate and interpret the results in photographic film.




Fig 1: Quick-blotsTM Mouse Tissues (Cat No.: 16-1001): Different Mouse tissue Lysates around 20  $\mu$ g per lane were resolved by SDS-PAGE, trasfered on to PVDF membrane and stained with amido black. Lanes: 1. Molecular weight marker, 2. Brain, 3. Heart, 4. Small Intestine, 5. Kidney, 6. Liver, 7. Lung, 8. Muscle, 9. Stomach, 10. Spleen, 11. Ovary, 12. Testis