

## 32-2731: PRDX6 Recombinant Protein

**Alternative Name :** Peroxiredoxin-6, Antioxidant protein 2, 1-Cys peroxiredoxin, Acidic calcium-independent phospholipase A2, Non-selenium glutathione peroxidase, 24 kDa protein, Liver 2D page spot 40, Red blood cells page spot 12, 1-Cys PRX, aiPLA2, NSGPx, PRDX6, AOP2,

### Description

Source : Escherichia Coli. Peroxiredoxin- 6 Human Recombinant fused with a 20 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 244 amino acids (1-224 a.a.) and having a molecular mass of 27.1kDa. The Peroxiredoxin-6 is purified by proprietary chromatographic techniques. Peroxiredoxin 6 (PRDX6) belongs to the thiol-specific antioxidant protein family. PRDX6 is a bifunctional enzyme with 2 distinct active sites. PRDX6 is involved in redox regulation of the cell and can reduce Hydrogen peroxide and short chain organic, fatty acid, and phospholipid hydroperoxides. PRDX6 may have a role in the regulation of phospholipid turnover as well as in protection against oxidative injury. Furthermore, PRDX6 eases the oxidative stress and TGF-beta-induced abnormalities of human trabecular meshwork cells. In addition, PRDX6 is necessary for blood vessel integrity in injured skin. At acidic pH, PRDX6 binds to reduced phospholipids, however at cytosolic pH PRDX6 binds only to phospholipids that are oxidized which is compatible with the role for PRDX6 in the repair of peroxidized cell membranes. Hydrogen peroxide-mediated hyperoxidation of PRDX6 induces cell cycle arrest at the G2/M transition via up-regulation of iPLA2 activity. Overexpression of PRDX6 is linked to oligodendroglioma.

### Product Info

**Amount :** 20 µg  
**Purification :** Greater than 95.0% as determined by SDS-PAGE.  
**Content :** The Peroxiredoxin-6 solution contains 20mM Tris-HCl buffer (pH8.0) and 20% Glycerol.  
**Storage condition :** Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.  
**Amino Acid :** MGSSHHHHHH SGLVPRGSH MPGGLLLGDV APNFEANTTV GRIRFHDFLG DSWGILFSHP  
RDFTPVCTTE LGRAAKLAPE FAKRNVKLI A LSIDSVEDHL A WSKDINAYN CEEPTEKLPF  
PIIDDRNREL AILLGMLDPA EKDEKGMPTV ARVVFVFGPD KKLKLSILYP ATTGRNFDEI  
LRVVISLQLT AEKRVATPVD WKDGD SVMVL PTIPEEEAKK LFPKGVFTKE LPSGKKYLR Y TPQP.

### Application Note

The specific activity was found to be approximately 95-120 pmole/min/Åµg.

