

32-2822: Streptokinase Recombinant Protein

Alternative Name : Streptokinase,SK.

Description

Source : Escherichia Coli. Streptokinase Recombinant produced in E.Coli is a non-glycosylated polypeptide chain containing 414 amino acids and having a molecular weight of 47.3kDa. The Streptokinase is purified by proprietary chromatographic techniques. Streptokinase is an extracellular metallo-enzyme produced by beta-haemolytic streptococcus and is used as an effective and cheap clot-dissolving medication in some cases of myocardial infarction (heart attack) and pulmonary embolism. It belongs to a group of medications known as fibrinolytics, and works by activating plasminogen through cleavage to produce plasmin.

Product Info

Amount : 750000, IU

Purification : Greater than 97.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Content : Lyophilized from a 0.2µm filtered concentrated (1mg/ml) solution in PBS, pH 7.4.

Storage condition : Lyophilized Streptokinase although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Streptokinase should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Amino Acid : IAGPEWLLDR PSVNNSQLVV SVAGTVEGTN QDISLKFFEI DLTSRPAHGG KTEQGLSPKS
KLFATDSGAM PHKLEKADLL KAIQEQLIAN VHSNDDYFEV IDFASDATIT DRNGKVYFAD
KDGSVTLPIQ PVQEFLKKGH VRVRPYKEKP VQNQAKSVDV EYTVQFTPLN PDDDFRPALK
DTKLLKTLAI GDTITSQELL AQAQSILNKN HPGYTIYERD SSIVTHDNDI FRTLPMQDE
FTYHVKNREQ AYRINKKSGL NEEINNTDLI SEKYYVLKKG EKPYPDFDRS HLKLFYIKYV
DVNTNELLKS EQLLTASERN LDFRDLYDPR DKAKLLYNNL DAFGIMDYTL TGKVEDNHDD
TNRITVYMG KRPEGENASY HLAYDKDRYT EEEREVYSYL RYTGTPIPDN PNDK.

Application Note

It is recommended to reconstitute the lyophilized Streptokinase in sterile 18M-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The specific biological activity measured by the ability of fibrin lysis in agarose plate was found to be 80000U/mg.

