

32-3065: NME2 Recombinant Protein

Alternative Name Nucleoside diphosphate kinase B,NDPK-B,NDPKB,NM23-H2,NM23B,EC 2.7.4.6,NDP kinase B,C-myc
:
purine-binding transcription factor PUF,NDK B,NME2,puf,MGC111212.

Description

Source : Escherichia Coli. NME2 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 152 amino acids (1-152 a.a.) and having a molecular mass of 17.2kDa. The NME2 is purified by conventional chromatography. NME2 takes an important part in the synthesis of nucleoside triphosphates other than ATP. NME2 negatively controls Rho activity by interacting with AKAP13/LBC. NME2 acts as a transcriptional activator of the MYC gene. NME2 binds DNA non-specifically. NME2 is a heterodimeric enzyme functioning as a nucleoside diphosphate kinase. NME1 and NME2 contain 152 amino acids, A and B polypeptide chains of the NM23 enzyme, respectively. NME2 is identical to the beta subunit of human erythrocyte NDP kinase. NDP kinases participate in the synthesis of nucleoside triphosphates, and NM23 is involved in the regulation of signal transduction by complexing with G proteins, causing activation/inactivation of developmental pathways.

Product Info

Amount : 25 µg
Purification : Greater than 95.0% as determined by SDS-PAGE.
Content : The NME2 protein solution contains 20mM Tris-HCl pH-8, 1mM DTT, and 10% glycerol.
Storage condition : NME2 although stable 4°C for 4 weeks, should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.
Amino Acid : MANLERTFIA IKPDGVQRGL VGEIIKRFEQ KGFRLVAMKF LRASEEHLKQ HYIDLKDRPF
FPGLVKYMNS GPVVAMVWEG LNVVKTGRVM LGETNPADSK PGTIRGDFCI QVGRNIIHGS
DSVKSAEKEI SLWFKPEELV DYKSCAHDWV YE.

