

## 32-3149: A2M Recombinant Protein

**Alternative Name :** Alpha-2-macroglobulin,Alpha-2-M,A2M,CPAMD5,FWP007,S863-7,alpha 2M,DKFZp779B086.

### Description

Source : Human Plasma. Human Alpha-2 Macroglobulin is a tetrameric glycoprotein, produced in Human plasma and having a molecular mass of 725 kDa. Alpha-2 Macroglobulin is a serine proteases inhibitor, which inhibits coagulation by inactivating thrombin and Kallikrein, it inhibits fibrinolysis by inactivating plasmin and involved in insulin transport. Alpha-2-Macroglobulin is a large plasma protein, which is produced by the liver, it's composed of 4 identical subunits bound together by -S-S- bonds. A2M is able to inactivate many kinds of proteinases (including serine-, cysteine-, aspartic- and metalloproteinases). A2M has a 35 amino acid 'bait' region in its structure. Proteinases bind and cleave the 'bait' region become bound to A2M. Macrophage receptors recognize the proteinase-A2M complex and clear it from the system. A2M binds to and removes MMP-2 and MMP-9 (active forms of the gelatinase) from the circulation using scavenger receptors on the phagocytes. The levels of Alpha-2-macroglobulin are increased in nephrotic syndrome which is a condition where the kidneys start to leak out some of the smaller blood proteins. Due to its large size, A2-macroglobulin is retained in the bloodstream. Increased production of all proteins causes A2-macroglobulin concentration to increase. Chronic renal failure might lead to amyloid by alpha-2-macroglobulin. A2M is raised in cirrhosis, pregnancy and diabetes.

### Product Info

|                            |   |
|----------------------------|---|
| <b>Amount :</b>            | 1 mg  |
| <b>Purification :</b>      | Greater than 96.0%.   |
| <b>Content :</b>           | Lyophilized from 0.02M TRIS, 0.15M NaCl and 0.1M Sucrose, pH 8.0.   |
| <b>Storage condition :</b> | Human Alpha-2 Macroglobulin although stable at room temperature for 3 weeks, should be stored between 2-8°C. Do not freeze! |

### Application Note

It is recommended to reconstitute the lyophilized A2M in sterile 18MΩ-cm H<sub>2</sub>O.

