

32-3158: ADFP Recombinant Protein

Alternative Name : Adipophilin, Adipose differentiation-related protein, ADRP, ADFP, MGC10598.

Description

Source : Escherichia Coli. ADFP Human Recombinant produced in E.Coli is a single, non-glycosylated, Polypeptide chain containing 444 amino acids and having a molecular mass of 49 kDa. The protein contains an extra 8 amino acid His tag at N-terminus. The ADFP amino acid sequence is identical to UniProtKB/Swiss-Prot entry Q99541 amino acids 4-437. The ADFP is purified by proprietary chromatographic techniques. ADFP is related with the globule surface membrane material. ADFP is a major constituent of the globule surface. Rise in mRNA levels is one of the initial indications of adipocyte differentiation. Mycobacterium leprae regulates ADFP expression to facilitate the accumulation of lipids within infected macrophages for intracellular survival. ADFP is expressed in lipid droplets of vitamin A-storing hepatic stellate cells and additionally in lipid droplets of steatotic hepatocytes. ADFP expression has a role in clear cell renal carcinoma differentiation. ADFP is a component of the lipid droplets in THP-1 cells.

Product Info

| | |
|----------------------------|---|
| Amount : | 10 µg |
| Purification : | Greater than 95% as determined by SDS PAGE. |
| Content : | Human ADFP was lyophilized from 0.5mg/ml solution containing 20mM Tris pH-7.5, and 20mM NaCl. |
| Storage condition : | Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C. |
| Amino Acid : | MKHHHHHHS VAVDPQPSVV TRVNLPLVS STYDLMSSAY LSTKDQYPYL KSVCEMAENG VKTITSVAMT SALPIIQKLE PQIAVANTYA CKGLDRIEER LPILNQPSTQ IVANAKGAVT GAKDAVTTTTV TGAKDSVAST ITGVMDKTKG AVTGSVEKTK SVVSGSINTV LGSRRMQLVS SGVENALTKS ELLVEQYLPL TEELEKEAK KVEGFDLVQK PSYYVRLGSL STKLHSRAYQ QALSRVKEAK QKSQQTISQL HSTVHLIEFA RKNVYSANQK IQDAQDKLYL SWVEWKRSIG YDDTDESHCA EHIESRTLAI ARNLTQQLQT TCHTLLSNIQ GVPQNIQDQA KHMGMAGDI YSVFRNAASF KEVSDSLLTS SKGQLQKMKE SLDDVMDYLV NNTPLNLVVG PFYPQLTESQ NAQDQGAEMD KSSQETQRSEHKTH. |

Application Note

Add deionized water to prepare a working stock solution of approximately 0.5mg/mL and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.

