

32-4373: Recombinant Bacterial Outer Membrane Protein-A

Alternative Name : Outer Membrane Protein-A, OmpA.

Description

Source : Escherichia Coli. The OmpA protein is one of the main outer-membrane proteins of a large array of Gram-negative bacteria such as A.salmonicida, Shigella dysenteriae and E.coli. OmpA's major physiological functions include maintenance of the structural integrity and morphology of the cells and porin activity, as well as a role in conjugation and bacteriophage binding. Achromogenic atypical Aeromonas salmonicida is the causative agent of goldfish ulcer disease. Virulence of this bacterium is associated with the production of a paracrystalline outer membrane A-layer protein. The species specific structural gene for the monomeric form of A-protein was cloned into a pET-3d plasmid in order to express and produce a recombinant form of the protein in E.coli BL21(DE3). The induced protein was isolated from inclusion bodies by a simple solubilization-renaturation procedure and purified by ion exchange chromatography on Q-Sepharose to over 95% pure monomeric protein. Recombinant A-protein was compared by biochemical, immunological and molecular methods with the A-protein isolated from atypical A.salmonicida bacterial cells by the glycine and the membrane extraction methods.

Product Info

Amount :	200 µg
Purification :	Greater than 98.0% as determined by:(a) Analysis by SEC-HPLC.(b) Analysis by SDS-PAGE.
Storage condition :	Lyophilized Bacterial Outer Membrane Protein-A although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon C between 2-7 days and for future use reconstituted OmpA should be stored at 4 below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please avoid freeze-thaw cycles.
Amino Acid :	mdvvispndn tfvttslasv tkqpvldfst aqqnltnfs evgdlknngf ivleiqgegq fndaeirqwl sngfwrppft gllvnpndhg nfansgevdn vrkffkiisd gtqtlvthti dsngkrlrla lasdveetin fadaevelkl nlanqafkl sgsqgtvalt agalwnasyt adpvatkplf klglfqlsl tnagkatalv segflknig danisatdfa itnvtnqti qrdkvnltit gdvsafkkda ngnlvnkaga sigwkaaadg qsatavlgag nmaggvqnal aafgtlyvaa dntvpvpavn fnvkaeiaggd sqatynyfkd eladfiltr dgmkfdtitt gttsanlihi rdvsnilpte ggkifvtite yadhaangrg egtvlvtrka lsvtlpsgga vtlkpadvaa dvgasitagr qarlvfevet nqgevavkks naegvdiqng trgtaplvdv tl.

Application Note

It is recommended to reconstitute the lyophilized OmpA in sterile 0.4% NaHCO₃, not less than 100µg/ml, which can then be further diluted to other aqueous solutions. The interaction of bacterial and recombinant A-layer protein with murine macrophages was directed at determining the effect of A-protein on intracellular events that occur in primed macrophages. This was accomplished by measuring the cytotoxic product produced by peritoneal macrophages when exposed to A-protein coated latex beads. Thioglycolate elicited macrophages exhibited a low level of activation (18% cytotoxicity) that was significantly increased (48% cytotoxicity) in the presence of latex beads. Coating of the latex beads with each of the three A-protein products resulted in an increase of cytotoxicity (mean +/- SEM) from 48% to 91%.

