

10-6007: Monoclonal Antibody to IKK alpha (Clone: ABM10G9)

| | |
|--------------------------------|---|
| Clonality : | Monoclonal |
| Clone Name : | ABM10G9 |
| Application : | FACS, WB |
| Reactivity : | Mouse, Human |
| Gene : | CHUK |
| Gene ID : | 1147 |
| Uniprot ID : | O15111 |
| Format : | Purified |
| Alternative Name : | Inhibitor of nuclear factor kappa-B kinase subunit alpha, I-kappa-B kinase alpha, Conserved helix-loop-helix ubiquitous kinase, Nuclear factor NF-kappa-B inhibitor kinase alpha, NFKB1A, Transcription factor 16 |
| Isotype : | Mouse IgG1, Kappa |
| Immunogen Information : | Full length recombinant protein of IKK alpha was used as the immunogen for this antibody. |

Product Info

| | |
|----------------------------|---|
| Amount : | 25 µg / 100 µg |
| Purification : | Protein G Chromatography |
| Content : | 25 µg in 50 µl / 100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic. |
| Storage condition : | Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles. |

Application Note

WB: 2-4 µg/ml, FACS: 0.5-1 µg/10⁶




Figure-1: Western blot analysis of IKK alpha. Anti-IKK alpha antibody (Clone: ABM10G9) was used at 2 µg/ml on HeLa, MCF7, U87, mSmall Intestine, PC3 and 293 lysates.




Figure-2: Intra cellular flow analysis of IKK alpha in Jurkat using 0.5 $\mu\text{g}/10^6$ cells of antibody (Clone: ABM10G9). Green represents isotype control; red represents anti-IKK alpha antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Figure-3: Intra cellular flow analysis of IKK alpha in 293 using 0.5 $\mu\text{g}/10^6$ cells of antibody (Clone: ABM10G9). Green represents isotype control; red represents anti-IKK alpha antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Figure-4: Intra cellular flow analysis of IKK alpha in HeLa using 0.5 $\mu\text{g}/10^6$ cells of antibody (Clone: ABM10G9). Green represents isotype control; red represents anti-IKK alpha antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Figure-5: Western blot analysis of IKK alpha. Anti-IKK alpha antibody (Clone: ABM10G9) was used at 2 $\mu\text{g}/\text{ml}$ on 3T3, Raw and EL-4 lysates.