

## 10-4136: Monoclonal Antibody to hCD95 (Clone: BR18 )

|                                |   |
|--------------------------------|---|
| <b>Clonality :</b>             | Monoclonal  |
| <b>Clone Name :</b>            | BR18  |
| <b>Application :</b>           | FACS  |
| <b>Reactivity :</b>            | Human   |
| <b>Gene :</b>                  | FAS   |
| <b>Gene ID :</b>               | 355   |
| <b>Uniprot ID :</b>            | P25445  |
| <b>Format :</b>                | Purified  |
| <b>Alternative Name :</b>      | Apo-1 antigen,apoptosis antigen 1,Apoptosis-mediating surface antigen FAS,APT1FASTM,Fas (TNF receptor superfamily,member 6), FASLG receptor |
| <b>Isotype :</b>               | Mouse IgG1 Kappa  |
| <b>Immunogen Information :</b> | A partial length Recombinant protein of human CD95 was taken as immunogen for this antibody.  |

### Product Info

|                            |   |
|----------------------------|---|
| <b>Amount :</b>            | 25 µg / 100 µg  |
| <b>Purification :</b>      | Protein G Chromatography  |
| <b>Content :</b>           | 25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.                |
| <b>Storage condition :</b> | Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles. |

### Application Note

FACS: 0.5-1 µg/10<sup>6</sup> cells




Fig-1: Cell Surface flow analysis of CD95 in PBMC (Lymphocytes) using 0.5 µg/10<sup>6</sup> cells. Green represents isotype control (ABEOMICS); red represents anti-hCD95 antibody (10-4136). Goat anti-mouse PE conjugated secondary antibody (ABEOMICS) was used.