

ABGENEX Pvt. Ltd., E-5, Infocity, KIIT Post Office, Tel: +91-674-2720712, +91-9437550560 Email: info@abgenex.com

Bhubaneswar, Odisha - 751024, INDIA

10-3003-NALE: Monoclonal Antibody to TLR9 (Clone: ABM1C51)

Clone Name: ABM1C51
Application: IHC,FACS,WB
Reactivity: Mouse,Human

 Gene :
 TLR9

 Gene ID :
 54106

 Uniprot ID :
 Q9NR96

Format: Azide Free, Purified

Alternative Name: TLR9,UNQ5798/PRO19605

Isotype: Mouse IgG1 Kappa

Immunogen Information: A partial length recombinant TLR9 protein (amino acids 100-290) was used as the immunogen for

the antibody.

Description

TLR9, a member of toll-like receptor family are central to the innate immunity by identifying pathogen associated molecular patterns (PAMPs). TLR9 identify unmethylated CpG dinucleotides present in bacterial DNA leading to NF-kB activation.

Product Info

Amount : 100 μg

Purification: Protein G Chromatography

Content : 25 μg in 50 μl/100 μg in 200 μl PBS containing No Azide, low endotoxin (0.1EU/1ug).

Storage condition:

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Western blot analysis: 2-4 μg/ml, FACS analysis: 0.5 μg/10⁶ cells, Immunohistochemical analysis: 5 μg/ml




Fig-1: Immunohistochemical analysis of TLR9 in human stomach tissue using TLR9 antibody (Clone: ABM1C51) at 5 μ g/ml.






Fig-2: Western blot analysis of TLR9. Anti- TLR9 antibody (Clone: ABM1C51) was used at 2 µg/ml on (1) Raji and (2) EL4 lysates.




Fig-3: Intracellular flow analysis of TLR9 in human PBMC (Lymphocytes) using 0.5 µg/10^6 cells of TLR9 antibody (Clone: ABM1C51). Green represents isotype control; red represents anti-TLR9 antibody. Goat anti-mouse PE conjugate was used as secondary.




Fig-4: Intracellular flow analysis of TLR9 in Raji cells using 0.5 μg/10⁶ cells of TLR9 antibody (Clone: ABM1C51). Green represents isotype control; red represents anti-TLR9 antibody. Goat anti-mouse PE conjugate was used as secondary antibody.




Fig-5: Immunohistochemical analysis of TLR9 in human Liver tissue using TLR9 antibody (Clone: ABM1C51) at 20 µg/ml.