

## 11-2003: Polyclonal Antibody to SARS-CoV-2 nucleocapsid Protein

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	ELISA, WB
<b>Format :</b>	Purified
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	A partial length recombinant coronavirus Nucleocapsid protein (amino acids 250-410) was used as the immunogen for this antibody.

### Product Info

<b>Amount :</b>	25 µg / 100 µg
<b>Purification :</b>	Protein A Chromatography
<b>Content :</b>	25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

Recommended dilutions: WB: 0.5-1 µg/ml. However, this need to be optimized based on the research applications.




Figure-1: Wells of a 96-microtiter plate were coated with 4µg/ml of SARS-Cov-2/nCov/COVID-19 Nucleocapsid recombinant protein (21-1003). The binding was detected by addition of different dilution of 11-2003 polyclonal antibody. The reactivity was detected by a HRP-conjugated goat-anti-rabbit polyclonal antibody.




Figure-2: Western Blot analysis of SARS-CoV-2 Nucleocapsid Protein: Anti- SARS-CoV-2 Nucleocapsid Protein (11-2003) was used at 2 µg/ml on (1) SARS-CoV-2 virus infected Vero Cell lysate and (2) Mock infected lysate.




Figure-3: Western Blot analysis of Nucleocapsid antibody: Anti- Nucleocapsid antibody (SARS-CoV-2) was used at 4 µg/ml on recombinant Nucleocapsid Protein (21-1003).