

ABGENEX Pvt. Ltd., E-5, Infocity, KIIT Post Office, Tel: +91-674-2720712, +91-9437550560 Email: info@abgenex.com

Bhubaneswar, Odisha - 751024, INDIA

32-1373: elL 1 beta Recombinant Protein

Alternative Name: Catabolin,Lymphocyte-activating factor (LAF),Endogenous Pyrogen (EP),Leukocyte Endogenous Mediator (LEM),Mononuclear Cell Factor (MCF),IL1F2,IL-1 beta.

Description

Source: Escherichia Coli. Recombinant IL 1 beta Equine produced in E.coli cells is a non-glycosylated, homodimeric protein containing 153 amino acid chain and having a molecular mass of 17.3kDa. The IL 1 beta is purified by proprietary chromatographic techniques. Interleukin-1b is produced by activated macrophages, IL-1B stimulates thymocyte proliferation by inducing il-2 release, b-cell maturation and proliferation, and fibroblast growth factor activity. IL1B proteins are involved in the inflammatory response, being identified as endogenous pyrogens, and are reported to stimulate the release of prostaglandin and collagenase from synovial cells.

Product Info

Amount: 10 µg

Purification: Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: The IL 1 beta was lyophilized from a 0.2µm filtered concentrated solution in PBS pH 7.4,

containing 0.1 % Tween-80.

Lyophilized IL 1 beta although stable at room temperature for 3 weeks, should be stored

Storage condition:

desiccated below -18°C. Upon reconstitution IL 1 beta should be stored at 4°C between 2-7 days

and for future use below -18°C.For long term storage it is recommended to add a carrier protein

(0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: AAMHSVNCRL RDIYHKSLVL SGACELQAVH LNGENTNQQV VFCMSFVQGE EETDKIPVAL

GLKEKNLYLS CGMKDGKPTL QLETVDPNTY PKRKMEKRFV FNKMEIKGNV EFESAMYPNW

YISTSQAEKS PVFLGNTRGG RDITDFIMEI TSA

Application Note

It is recommended to reconstitute the lyophilized IL 1 beta in sterile distilled H2O not less than $100\text{Å}\mu\text{g/ml}$, which can then be further diluted to other aqueous solutions. The ED50 as determined by a cell proliferation assay using murine D10S cells is less than 20 pg/ml, corresponding to a specific activity of > $5.0 \times 107 \text{ IU/mg}$.

