

ABGENEX Pvt. Ltd., E-5, Infocity, KIIT Post Office, Tel: +91-674-2720712, +91-9437550560 Email: info@abgenex.com

Bhubaneswar, Odisha - 751024, INDIA

## 32-1522: IL36RN Recombinant Protein

Alternative Name :

Interleukin-36 receptor antagonist protein,FIL1 delta,IL-1-related protein 3,IL-1RP3,Interleukin-1 HY1,IL-1HY1,Interleukin-1 delta,IL-1 delta,Interleukin-1 family member 5,IL-1F5,Interleukin-1 receptor

antagonist homolog 1,IL-1ra homolog 1

## **Description**

Source: Escherichia Coli. IL1F5 Human Recombinant produced in E.Coli is a single, non-glycosylated, Polypeptide chain containing 155 amino acids and having a molecular mass of 17kDa.The IL1F5 is purified by proprietary chromatographic techniques. Human interleukin family 1, member 5 (IL-1F5 / FIL1-delta) belongs to the interleukin 1 cytokine family. IL1F5 is expressed by a variety of cells including monocytes, Bcells, dendritic cells/Langerhans cells, keratinocytes, and gastric fundus Parietal and Chief cells. IL1F5 is an antagonist of IL1F9; however IL1F5 activity related to receptor binding remains unclear. Human and mouse IL1F5 share 90% amino acid sequence identity.

## **Product Info**

Amount: 25 µg

Purification: Greater than 95.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content: IL1F5 was lyophilized after extensive dialysis against 20mM Phosphate buffer, pH7.4.

ictro was hyphilized after extensive dialysis against 2011iiii rhospitate buffer, pri7.4.

Lyophilized IL1F5 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL1F5 should be stored at 4°C between 2-7 days and for future

use below -18°C.Please prevent freeze-thaw cycles.

Amino Acid: The sequence of the first five N-terminal amino acids was determined and was found to be Met-Val-

Leu-Ser-Gly.

## **Application Note**

It is recommended to quick spin followed by reconstitution of IL1F5 in PBS to a concentration no less than 100 Âμg/ml, which can then be further diluted to other aqueous solutions. As measured by its binding ability in a functional ELISA, immobilized IL1F5 at 1 Âμg/ml (100 Âμl/well) can bind rHuIL-1 Rrp2/Fc Chimera with a linear range of 0.15- 5 Âμg/ml.

