

ABGENEX Pvt. Ltd., E-5, Infocity, KIIT Post Office, Tel: +91-674-2720712, +91-9437550560 Email: info@abgenex.com

Bhubaneswar, Odisha - 751024, INDIA

## 32-1529: IL36B Recombinant Protein

Alternative Name Interleukin 36 beta,interleukin 1 family member 8 (eta),Interleukin-1 homolog 2,IL1F8 (Canonical product IL-1F8a),IL-1F8 (FIL1-eta),Interleukin-1 Superfamily e,IL1H2,MGC126880,MGC126882.

## **Description**

Source: Escherichia Coli. IL36B Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 157 amino acids and having a molecular mass of 17.7kDa.The IL36B is purified by proprietary chromatographic techniques. Human IL-36b belongs to the IL-1 family that includes IL-1b, IL-1a, IL-1ra, IL-18, IL-36ra (IL1F5), IL-36b (IL1F8), IL-36g (IL1F9), IL-37 (IL1F7) and IL-38 (IL-1F10). The IL-1 family members display a 12 b-strand, b-trefoil configuration, and are thought to have ascended from a mutual ancestral gene. IL-36 beta is known to be actively secreted. Cells expressing IL-36 beta include resting and activated monocytes and B cells. The receptor for IL-36 beta is a blend of IL-1 Rrp2 and IL-1 RAcP. Recombinant IL-36 beta stimulates processes involving NF-kB and MAPK in an IL-1 Rrp2-dependent manner.

## **Product Info**

Amount: 10 µg

**Purification:** Greater than 95.0% as determined by SDS-PAGE.

Content: Lyophilized from a 0.2µm filtered concentrated solution in 1xPBS, pH 7.4.

Lyophilized IL36B Human although stable at room temperature for 3 weeks, should be stored

Storage condition: Lyophilized IL36B Human although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL36B should be stored at 4°C between 2-7 days and

for future use below -18°C. Please prevent freeze-thaw cycles.

Amino Acid: MNPQREAAPK SYAIRDSRQM VWVLSGNSLI AAPLSRSIKP VTLHLIACRD TEFSDKEKGN

MVYLGIKGKD LCLFCAEIQG KPTLQLKEKN IMDLYVEKKA QKPFLFFHNK EGSTSVFQSV

SYPGWFIATS TTSGQPIFLT KERGITNNTN FYLDSVE

## **Application Note**

It is recommended to reconstitute the lyophilized IL36B in sterile 18M-cm H2O not less than 100Âμg/ml, which can then be further diluted to other aqueous solutions. Measured by its binding ability in a functional ELISA to bind recombinant human IL-1 Rrp2 Fc Chimera.

