

## 32-1530: IL36B 153 a.a. Recombinant Protein

**Alternative Name** Interleukin 36 beta,interleukin 1 family member 8 (eta),Interleukin-1 homolog 2,IL1F8 (Canonical product : IL-1F8a),IL-1F8 (FIL1-eta),Interleukin-1 Superfamily e,IL1H2,MGC126880,MGC126882.

### Description

Source : Escherichia Coli. IL36B 153 a.a. Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 153 amino acids (5-157a.a.) and having a molecular mass of 17.2kDa.The IL36B 153 a.a. Human is purified by proprietary chromatographic techniques. Human IL-36b belongs to the IL-1 family that includes IL-1b, IL-1a, IL-1ra, IL-18, IL-36ra (IL1F5), IL-36b (IL1F8), IL-36g (IL1F9), IL-37 (IL1F7) and IL-38 (IL-1F10). The IL-1 family members display a 12 b-strand, b-trefoil configuration, and are thought to have ascended from a mutual ancestral gene. IL-36 beta is known to be actively secreted. Cells expressing IL-36 beta include resting and activated monocytes and B cells. The receptor for IL-36 beta is a blend of IL-1 Rrp2 and IL-1 RAcP. Recombinant IL-36 beta stimulates processes involving NF-kB and MAPK in an IL-1 Rrp2-dependent manner.

### Product Info

<b>Amount :</b>	10 µg
<b>Purification :</b>	Greater than 97.0% as determined by SDS-PAGE and HPLC analyses.
<b>Content :</b>	Lyophilized from a 0.2µm filtered concentrated solution in 1xPBS, pH 7.4.
<b>Storage condition :</b>	Lyophilized IL36B 153 a.a. Human although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL36B 153 a.a. Human should be stored at 4°C between 2-7 days and for future use below -18°C.Please prevent freeze-thaw cycles.
<b>Amino Acid :</b>	REAAPKSYAI RDSRQMVVWL SGNSLIAAPL SRSIKPVT LH LIACRDTEFS DKEKGNMVYL GIKGDLCFL CAEIQGKPTL QLKEKNIMDL YVEKKAQKPF LFFHNKEGST SVFQSVSYPG WFIATSTTSG QPIFLTKERG ITNNTNFYLD SVE

### Application Note

It is recommended to reconstitute the lyophilized IL36B 153 a.a. Human in sterile 18M-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions. Fully biologically active when compared to standard. The ED50 as measured by its ability to induce IL-8 secretion in human preadipocytes is less than 10ng/ml, corresponding to a specific activity of 10IU/mg.

