

32-1867: CXCL17 Recombinant Protein

Alternative Name VEGF coregulated chemokine 1,C-X-C motif chemokine 17,Dendritic cell and monocyte chemokine-like protein,DMC,CXCL17,VCC1,Dcip1,VCC-1,UNQ473.

Description

Source : Escherichia Coli. CXCL17 Human Recombinant produced in E.coli is a single, non-glycosylated polypeptide chain containing 98 amino acids and having a molecular mass of 11.5kDa. The CXCL17 is purified by proprietary chromatographic techniques. Dendritic cell and monocyte chemokine like protein (DMC/CXCL17/VEGF-correlated chemokine 1/VCC1), is a secreted molecule with a size and predicted 3-dimensional folding pattern similar to that of chemokines CXCL8/IL8 and CXCL14/BRAK. CXCL17 is constitutively generated by airway and intestinal epithelium. CXCL17 induces the chemotaxis of quiescent, but not LPS-activated peripheral blood monocytes and dendritic cells, and it also binds these cells specifically. The expression of CXCL17 is increased in endothelial cells when they are induced to form tubes in vitro. CXCL17, CXCL1/GRO and CXCL8/IL8 which have roles in angiogenesis, show significantly correlated expression with that of VEGF in primary lung, breast and esophageal tumors. Therefore, CXCL17 is suggested to have a role in tumor angiogenesis. The mature Rat CXCL17 shares 82%, 71% amino acid sequence identity with mouse, human CXCL17, respectively.

Product Info

Amount :	25 µg
Purification :	Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content :	CXCL17 protein was lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4, containing 3% Trehalose.
Storage condition :	Lyophilized CXCL17 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CXCL17 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.
Amino Acid :	SSLNPGVARG HRDRGQASRR WLQEGGQECE CKDWFLRAPR RKFMTVSGLP KKQCPCDHFK GNVKTRHQR HHRKPNKHSR ACQQFLKQCQ LRSFALPL.

Application Note

It is recommended to reconstitute the lyophilized CXCL17 in sterile 18M-cm H₂O not less than 100Åµg/ml, which can then be further diluted to other aqueous solutions. The ED₅₀ as determined by its ability to induce VEGF expression using murine endothelial cells is less than 5.0Åµg/ml, corresponding to a specific activity of > 200IU/mg.

