

32-1941: MCP 4 Recombinant Protein

Alternative Name : Small inducible cytokine A13,CCL13,Monocyte chemotactic protein 4,MCP-4,Monocyte chemoattractant protein 4,CK-beta-10,NCC-1,chemokine (C-C motif) ligand 13,NCC1,CKb10,SCYL1,SCYA13,MGC17134.

Description

Source : Escherichia Coli. Monocyte Chemotactic Protein-4 Human Recombinant produced in E.Coli is a non-glycosylated, Polypeptide chain containing 75 amino acids and having a molecular mass of 8.6 kDa. The MCP-4 is purified by proprietary chromatographic techniques. Chemokine (C-C motif) ligand 13 (CCL13 / MCP-4) is a small cytokine belonging to the CC chemokine family. The MCP-4 gene is located on human chromosome 17 within a large cluster of other CC chemokines. MCP-4 induces chemotaxis in monocytes, eosinophils, T lymphocytes, and basophils by binding cell surface G-protein linked chemokine receptors such as CCR2, CCR3 and CCR5. Activity of the MCP-4 chemokine has been implicated in allergic reactions such as asthma. MCP-4 can be induced by the inflammatory cytokines interleukin-1 and TNF- α .

Product Info

Amount : 10 μ g
Purification : Greater than 96.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.
Content : The protein was lyophilized from a concentrated (1mg/ml) sterile solution in 20mM PB, pH 7.4, 130mM NaCl.
Storage condition : Lyophilized MCP-4 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution MCP-4 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.
Amino Acid : QPDALNVPSTCCFTFSSKKISLQRLKSYVITTSRCPQKAVIFRTKLGKEICADPKEKWVQNYMKHLGRKAHTLKT.

Application Note

It is recommended to reconstitute the lyophilized MCP-4 in sterile 18M-cm H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions. The specific activity as determined by the ability of MCP-4 to chemoattract human monocytes at 10-100ng/ml, corresponding to a Specific Activity of 10,000-100,000 units/mg.

