

ABGENEX Pvt. Ltd., E-5, Infocity, KIIT Post Office, Tel: +91-674-2720712, +91-9437550560 Email: info@abgenex.com

Bhubaneswar, Odisha - 751024, INDIA

## 32-1982: mPF 4 Recombinant Protein

Alternative Name: CXCL4,PF-4,PF4,Iroplact,Oncostatin-A,SCYB4,MGC138298.

## **Description**

Source: Escherichia Coli. CXCL4 Mouse Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 76 amino acids and having a molecular mass of 8.2kDa. Platelet factor-4 is a 70-amino acid protein that is released from the alpha-granules of activated platelets and binds with high affinity to heparin. Its major physiologic role appears to be neutralization of heparin-like molecules on the endothelial surface of blood vessels, thereby inhibiting local antithrombin III activity and promoting coagulation. As a strong chemoattractant for neutrophils and fibroblasts, PF4 probably has a role in inflammation and wound repair. Oncostatin-A is a member of the CXC chemocinfamily. Human PF4 is used for the proof of heparin-induced thrombocytopenia. Furthermore it is used as an inhibitor in the angiogenesis during tumor therapy.

## **Product Info**

Amount: 20 µg

Purification: Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Content:

The Mouse CXCL4 protein was lyophilized from a 0.2µm filtered concentrated solution in 20mM

PB, pH 7.4 and 1.0M NaCl.

 $Lyophilized \ CXCL4 \ Mouse \ Recombinant \ although \ stable \ at \ room \ temperature \ for \ 3 \ weeks, \ should$ 

Storage condition:

be stored desiccated below -18°C. Upon reconstitution Mouse CXCL4 should be stored at 4°C

between 2-7 days and for future use below -18°C.For long term storage it is recommended to add

a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Amino Acid: VTSAGPEESD GDLSCVCVKT ISSGIHLKHI TSLEVIKAGR HCAVPQLIAT LKNGRKICLD

RQAPLYKKVI KKILES.

## **Application Note**

It is recommended to reconstitute the lyophilized CXCL4 in sterile 18MΩ-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

