

36-1939: Monoclonal Antibody to CD44 / HCAM Std.(Clone : SPM544)

Clonality :	Monoclonal
Clone Name :	SPM544
Application :	IF,Functional Assay,WB,IHC
Reactivity :	Human
Gene :	CD44
Gene ID :	960
Uniprot ID :	P16070
Format :	Purified
Alternative Name :	CD44,LHR,MDU2,MDU3,MIC4
Isotype :	Mouse IgG2a, kappa
Immunogen Information :	Stimulated human leukocytes

Description


Recognizes a cell surface glycoprotein of 80-95kDa (CD44) on lymphocytes, monocytes, and granulocytes. Its epitope is resistant to digestion by trypsin and chymotrypsin. The CD44 family of glycoproteins exists in a number of variant isoforms, the most common being the standard 85-95kDa or hematopoietic variant (CD44s). Higher molecular weight isoforms are described in epithelial cells (CD44v), which are believed to function in intercellular adhesion and stromal binding. CD44 immunostaining is commonly used for the discrimination of urothelial transitional cell carcinoma in-situ from non-neoplastic changes in the urothelium.

Product Info

Amount :	100 µg
Purification :	Affinity Chromatography
Content :	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Immunofluorescence (1-2ug/ml); Functional Studies (Order Ab without BSA & Azide); Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (0.25-0.5ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Formalin-fixed, paraffin-embedded human Tonsil stained with CD44 Monoclonal Antibody (SPM544)