

36-1970: Monoclonal Antibody to CD68 (Macrophage Marker)(Clone : CD68/G2)

| | |
|--------------------------------|--------------------------------|
| Clonality : | Monoclonal |
| Clone Name : | CD68/G2 |
| Application : | IHC |
| Reactivity : | Human |
| Gene : | CD68 |
| Gene ID : | 968 |
| Uniprot ID : | P34810 |
| Format : | Purified |
| Alternative Name : | CD68 |
| Isotype : | Mouse IgG1, kappa |
| Immunogen Information : | Recombinant human CD68 protein |

Description


This antibody recognizes a glycoprotein of 110kDa, which is identified as CD68. It is important for identifying macrophages in tissue sections. It stains macrophages in a wide variety of human tissues, including Kupffer cells and macrophages in the red pulp of the spleen, in lamina propria of the gut, in lung alveoli, and in bone marrow. It reacts with myeloid precursors and peripheral blood granulocytes. It also reacts with plasmacytoid T cells, which are supposed to be of monocyte/macrophage origin. It shows strong granular cytoplasmic staining of chronic and acute myeloid leukemia and also reacts with rare cases of true histiocytic neoplasia. Lymphomas are negative or show few granules.

Product Info

| | |
|----------------------------|---|
| Amount : | 100 µg |
| Purification : | Affinity Chromatography |
| Content : | 100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic. |
| Storage condition : | Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles. |

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Formalin-fixed, paraffin-embedded human Histiocytoma stained with CD68 Monoclonal Antibody (CD68/G2).