

ABGENEX Pvt. Ltd., E-5, Infocity, KIIT Post Office, Tel: +91-674-2720712, +91-9437550560 Email: info@abgenex.com Bhubaneswar, Odisha - 751024, INDIA

10-3010: Monoclonal Antibody to TLR4 (Clone: ABM19C4)

Clonality: Monoclonal ABM19C4 Clone Name: Application: IHC.FACS.IF Reactivity: Human Gene: TLR4 7099 Gene ID: **Uniprot ID:** O00206 Format: Purified

Isotype: Mouse IgG2b Kappa

Immunogen Information: A partial length recombinant TLR4 protein (amino acids 45-320) was used as the immunogen for

this antibody.

TLR4

Description

Alternative Name:

TLR4 (Toll-like receptor-4) is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. It primarily mediates cellular signaling induced by Gram?negative bacteria. TLR4 is highly specific for LPS and it is associated with CD14, a co-receptor for LPS. The downstream signaling pathway of TLR4 leads to the activation of NF-kappaB through myeloid differentiation protein (MyD88) and IL-1 receptor-associated kinase in various cell types. Keratinocytes express TLR4 at the mRNA and protein levels whereas TLR4 are also present in the normal human epidermis in vivo and their expression is regulated by microbial components. TLR4 is expressed on CD4+ T cells as well, the functional significance of which is unclear.

Product Info

Amount : $25 \mu g / 100 \mu g$

Purification: Protein G Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition:

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Immunohistochemical analysis: 5 µg/ml; FACS analysis: 2-4 µg/10^6



ABGENEX Pvt. Ltd., E-5, Infocity, KIIT Post Office, Tel: +91-674-2720712, +91-9437550560 Email: info@abgenex.com

Bhubaneswar, Odisha - 751024, INDIA




Fig-1 : Immunohistochemical analysis of TLR4 in human skin tissue using TLR4 antibody (Clone: ABM19C4) at 5 μ g/ml.




Fig-2:Intracellular flow analysis of hTLR4 antibody in TLR4 Transfected Cell line using 2 μ g/10^6 cells of hTLR4 antibody (Clone: ABM19C4). Green represents isotype control; red represents anti-hTLR4 antibody. Goat anti-mouse FITC conjugate was used as secondary antibody.