

20-1093: Polyclonal antibody to TRAF-1

| | |
|--------------------------------|--|
| Clonality : | Polyclonal |
| Application : | IP,IHC,WB |
| Reactivity : | Human |
| Gene : | TRAF1 |
| Gene ID : | 7185 |
| Uniprot ID : | Q13077 |
| Format : | Sera |
| Alternative Name : | TRAF1,EBI6 |
| Isotype : | Rabbit IgG |
| Immunogen Information : | A synthetic peptide of human TRAF-1 (amino acids 1-22 MASSSGSSPRPAPDENEFPFG) was used as immunogen for this antibody |

Description

This antibody recognizes TRAF1. TRAF1 is a member of TRAF (TNF receptor-associated factor) adapter proteins composed 416 amino acids. The members of this family link a wide variety of cell surface receptors to diverse signaling cascades leading to the activation of NF- κ B and mitogen-activated protein kinases. TRAFs are major signal transducers for both the TNF and IL-1/TLR receptor superfamilies and collectively play important functions in both adaptive and innate immunity. TRAFs also interact with a variety of proteins that regulate receptor-induced cell death or survival, and TRAF-mediated signaling can promote cell survival or interfere with death receptor-induced apoptosis.

Product Info

| | |
|----------------------------|---|
| Amount : | 50 μ l |
| Content : | 50 μ l sera |
| Storage condition : | Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles. |

Application Note

WB: 1:1000-1:2000, IHC (paraffin): 1:1000-1:5000, IHC (frozen): Users should optimize, IP: 1:50-1:200




Fig:1 Western blot analysis of TRAF1 in normal human tissues using 20-1093 at 1:2000. TRAF1 is observed at 50 kDa. Additional bands of lower molecular weight were seen in some cases, and may represent TRAF1 degradation fragments.




Fig:2 Formalin-fixed, paraffin-embedded human lymphoma tissue section stained for TRAF1 expression using 20-1093 at 1:2000. Hematoxylin-eosin counterstain. A1 is a higher magnification of A.




Fig:3 Formalin-fixed, paraffin-embedded human tissue sections stained for TRAF1 expression using 20-1093 at 1:2000. Hematoxylin-eosin counterstain. A, A1: Formalin-fixed, paraffin-embedded human lymphoma tissue section stained for TRAF1 expression using 20-1093 at 1:2000. Hematoxylin-eosin counterstain. A, Hodgkin's Disease, the Reed-Sternberg cells are positive for TRAF1 expression. B, lymph node, dendritic-like cells are positive for TRAF1 expression. A1 and B1 are higher magnifications from A and B, respectively.