

ABGENEX Pvt. Ltd., E-5, Infocity, KIIT Post Office, Tel: +91-674-2720712, +91-9437550560 Email: info@abgenex.com

Bhubaneswar, Odisha - 751024, INDIA

## 20-1087: Polyclonal antibody to SMAC/Diablo

Clonality: Polyclonal IP,IHC,WB Application: Reactivity: Human DIABLO Gene: Gene ID: 56616 **Q9NR28 Uniprot ID:** Format: Sera

**Alternative Name:** DIABLO, SMAC Isotype: Rabbit IgG

A synthetic peptide of SMAC/Diablo (amino acids 56-73 AVPIAQKSEPHSLSSEAL) was used as Immunogen Information:

immunogen for this antibody

## **Description**

Smac/DIABLO (second mitochondrial-derived activator of caspase/direct IAP binding protein with low pI) is encoded by the nucleus and resides as a precursor within the intermembranous space of the mitochondria. During apoptosis, Smac/DIABLO is proteolytically cleaved and released from the mitochondria into the cytosol, where it promotes cell death by interacting with IAPs and inhibiting their anti-apoptotic function. In turn, certain IAPs may degrade Smac/DIABLO and block its cell death function. Smac/DIABLO is expressed in a wide range of normal tissues and some tumor cells lines, and its over-expression in tumor cell lines can trigger both extrinsic and intrinsic apoptosis pathways.

## **Product Info**

Amount: 50 µl Content: 50 µl sera

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid Storage condition:

repeated freeze and thaw cycles.

## **Application Note**

WB: 1:1000-1:2000, IHC (paraffin): 1:1000-1:5000, IHC (frozen): Users should optimize, IP: 1:50-1:200



Fig:1 Formalin-fixed, paraffin-embedded tissue cores from a human breast cancer microarray stained for Smac/Diablo using 20-1087 at 1:2000. Hematoxylin-eosin counterstain. Differential Smac/DIABLO staining is seen between patient samples.