

11-316: Alkaline phosphatase conjugated Goat anti Rabbit IgG (H+L)

Clonality : Polyclonal
Application : ELISA, WB
Reactivity : Rabbit
Isotype : Goat IgG

Description

Secondary antibodies are antibodies that bind to primary antibodies or antibody fragments. They are typically labeled with probes that make them useful for detection, purification or cell sorting applications. Secondary antibodies may be polyclonal or monoclonal, and are available with specificity for whole Ig molecules or antibody fragments such as the Fc or Fab regions. Secondary antibodies are selected according to the source of the primary antibody, the class of the primary antibody (e.g., IgG or IgM), and the kind of label which is preferred. Identifying the optimal secondary antibody is normally done through trial and error method. Secondary antibodies are used in all types of immunoassays, most often in Western blot, immunohistochemistry, and immunocytochemistry, and occasionally in immunoprecipitation. Basic research, clinical analysis, and disease diagnosis also use secondary antibodies in ELISA and flow cytometry assays. They are also useful for cell sorting, fluorescence activated cell sorting, FACS.

Product Info

Amount : 25 µl / 100 µl
Purification : Affinity Chromatography
Content : Antibody is suspended in 0.01 M Tris-HCl, 0.25 M NaCl, pH 8.0
Storage condition : Store the antibody at 4°C, stable for 6 months.

Application Note

ELISA / Western blot analysis: 1:5,000-1:20,000

